

# Glueball-Meson Mixing

Vicente Vento

Departamento de Física Teórica and Instituto de Física Corpuscular, Universidad de Valencia -  
Consejo Superior de Investigaciones Científicas, 46100 Burjassot (València), Spain,  
Email: vicente.vento@uv.es.

## Abstract

Calculations in unquenched QCD for the scalar glueball spectrum have confirmed previous results of Gluodynamics finding a glueball at  $\sim 1750$  MeV. I analyze the implications of this discovery from the point of view of glueball - meson mixing at the light of the experimental scalar spectrum.

Pacs: 12.38.-t, 12.38.Aw, 14.70.-e, 14.80.-j

Keywords: meson, glueball, mixing.

# 1 Introduction

Glueballs have not been an easy subject to study due to the lack of phenomenological support and therefore much debate has been associated with their properties [1]. I center the discussion here at the consequences of the spectrum obtained by lattice QCD where the results seem to be converging. In quenched QCD the masses of the scalar glueballs appear large  $\geq 1700$  MeV [2–5], a result which has been confirmed by unquenched calculations [6]. .

Several glueball-meson mixing scenarios have been discussed in the literature using either lattice calculations or phenomenology [2, 3, 7–9]. I implement a combination of lattice results and phenomenology to study the implications of recent lattice results in the possible mixing scenarios in the scalar sector.

## 2 The scalar spectrum

Lattice QCD provides us with a value for the mass of the  $0^{++}$  glueball states as shown in Table 1.

$J^{PC}$	Mass MeV		
	Unquenched	Quenched	
	Gl	Mp	Ky
$0^{++}$	1795(60)	1730(50)(80)	1710(50)(80)
$0^{++}$	3760(240)	2670(180)(130)	

Table 1: Glueball masses with  $J^{PC}$  assignments. The column Gl reports the results of the unquenched QCD calculation by Gregory et al. [6], the columns Mp and Ky show the data from Morningstar and Peardon [4] and Chen et al. [5] respectively.

The three calculations give a similar mass for their lowest state for which I take the mean  $1743 \pm 42$  MeV in my analysis.

In Table 2, I show the experimental scalar spectrum, namely that of the particles labelled  $f_0$ .

$J^{PC}$	Name	Mass MeV	Width MeV	Comment
$0^{++}$	$f_0(500)$	400-550	400-700	
$0^{++}$	$f_0(980)$	$990 \pm 20$	40-100	
$0^{++}$	$f_0(1370)$	1200-1500	200-500	
$0^{++}$	$f_0(1500)$	$1505 \pm 6$	$109 \pm 7$	
$0^{++}$	$f_0(1710)$	$1720 \pm 6$	$135 \pm 8$	
$0^{++}$	$f_0(2020)$	$1992 \pm 16$	$442 \pm 60$	needs confirmation
$0^{++}$	$f_0(2100)$	$2013 \pm 8$	$209 \pm 19$	needs confirmation
$0^{++}$	$f_0(2200)$	$2189 \pm 13$	$238 \pm 50$	needs confirmation

Table 2: Scalar particles appearing in the PDG’s particle listings [10].

The three heaviest states have not been confirmed and some authors also question the existence of the  $f_0(1370)$ .

An observation at the light of the experimental spectrum is that the excited glueballs obtained in QCD calculations are very high in mass and therefore I do not expect them to mix with the mesons. Thus the lattice calculations and the observed scalar spectrum lead to a scenario of one glueball amidst several scalar mesons in the range between  $1 - 2$  GeV.

### 3 Glueball-Meson Mixing

The most naive way of implementing a mixing scenario is by assuming a hamiltonian which is not diagonal in the unmixed states. Thus there are two ingredients to this hamiltonian, the unmixed masses and the off-diagonal mixing parameters.

The calculations thus far have led to different realizations of mixing. Some obtain that the  $f_0(1710)$  is mostly glueball with small admixtures of  $q\bar{q}$  states [2, 3, 9]. Others claim that the  $f_0(1500)$  is mostly glueball, with small admixtures of  $q\bar{q}$  states [7, 8].




Figure 1: The largest eigenvalue of the studied analog of a three state mixing matrix is shown as a function of the relevant parameters in Cardano's formulas. The values used for the diagonal elements were 1750, 1500, 1300 close to the physical  $f_0$  masses, and the mixing parameters  $b, c$  range from  $0 \rightarrow 1000$ , and  $a$  from  $-1000 \rightarrow 1000$  in discrete steps.

I analyze next which scenarios could arise in a full QCD calculation. The idea is to construct a hamiltonian which is not diagonal in the unmixed lattice states and diagonalize it to obtain the physical states. The decisive input is the glueball mass as given by lattice QCD. I assume that the rest of the particles in the unmixed Fock space are scalar mesons. Since the scalar meson spectrum of lattice QCD is not well known I will be guided by phenomenology.

For the off-diagonal matrix elements I shall use perturbation theory and  $1/N_c$  arguments. With these assumptions the matrix elements are of the form  $\sim | \langle \Psi_n | H_{non-diagonal} | \Psi_m \rangle |^2 / (E_n - E_m)$ , therefore they depend on the inverse of the decay coupling  $f_{meson}^2 \sim N_c$ ,  $f_{glueball}^2 \sim N_c^2$ , and on the inverse of the energy difference of the two levels. For almost degenerate unmixed states, the

mixing parameters might be larger than for states of quite different masses. Thus the hamiltonian matrix might look for a three state mixing as,

$$\begin{pmatrix} m_{s1} & a & b \\ a & m_{s2} & c \\ b & c & m_g \end{pmatrix}, \quad (1)$$

where,  $m_{s1}$  and  $m_{s2}$  represent the unmixed scalar meson masses, and  $m_g$  the unmixed glueball mass. The diagonal matrix elements I take motivated by phenomenology and their values incorporate all higher order corrections,  $\sim 1 + O(N_c^{-2}) + \dots$  for the glueballs, and  $\sim 1 + O(N_c^{-1}) + \dots$  for the mesons. The mixing matrix elements are  $\sim O(N_c^{-2})$  for meson-glueball states since  $\Delta E \sim 1/N_c$ , but they are also  $\sim O(N_c^{-2})$  for the meson-meson states since  $\Delta E \sim O(1)$  for them. Thus  $a, b$  and  $c$  will be of the same order of magnitude, i.e.  $\leq 250$  MeV.

It is trivial to show that in a two by two symmetric mass mixing matrix the the unmixed mass of the heavy state will increase after non trivial diagonalisation. For a three state mass mixing matrix it is not so trivial to show that the heaviest state mass also increases after non trivial diagonalisation. This can be done using Cardano's formulas, as shown in Fig. 1. Cardano's formulas depend only of one sign,  $sign(abc)$ , therefore one reaches all the desired values by only letting one of the parameters become negative. I have used as diagonal elements numbers close to the actual  $f_0$  masses and the off diagonal elements have taken values far beyond my expectation for the physical mixing parameter values.




Figure 2: Left: Eigenvalues of the mixing matrix for the glueball and two meson states. The masses and their errors are from PDG [10]. Right: The probability distribution for the  $f_0(1710)$  of the unmixed states as a function of  $c$ , having fixed  $a = b = 50$  MeV. The dotted line represents the glueball with unmixed mass (1670 MeV), the full line the heavy meson with unmixed mass (1530 MeV), the dashed line the light meson with unmixed mass ( $\sim 1380$  MeV). The error bars in the probability curves are just to guide the eye.

The QCD lattice mass value for the scalar glueball is very high and the error relatively small if the three values shown in Table 1 are averaged with errors in quadrature  $m_g = 1743 \pm 42$  MeV. One can still argue, given the errors in the lattice calculation, that the glueball mass could be below, but certainly not much below, the experimental  $f_0(1710)$ . Therefore, I foresee two scenarios: i)

the unmixed QCD mass value is below the  $f_0(1710)$  but close to it, and ii) the unmixed glueball mass value is above the  $f_0(1710)$  in agreement with the lattice QCD average value. In the latter case I will analyze two cases depending on the unmixed spectrum, one will be associated with weak mixing, while the other with strong mixing.




Figure 3: Left: Eigenvalues of the mixing matrix for the glueball and two meson states. The masses and their errors are from PDG [10]. Right: The probability distribution for the  $f_0(1710)$  of the unmixed states as a function of  $c$ . The other mixing parameters have been fixed to  $a = 100$  MeV and  $b = 100$ . The dotted line represents the unmixed glueball (1750), the dashed line represents the light unmixed meson (1550), the full line represents the heavy unmixed meson (1900)

The first scenario requires very small mixing, as can be seen in Fig.2 (left). The unmixed values were chosen  $m_{s1} = 1380$  MeV,  $m_{s2} = 1530$  MeV and  $m_g = 1670$  MeV, the latter two standard deviations below the central lattice value. The mixing parameters that fit the data are small  $a = b = 50$  MeV and  $c = 70$  MeV. Note that in this case one can also trivially construct a two state weak mixing scenario, with the outcome that the  $f_0(1710)$  is again mostly glueball, the  $f_0(1500)$  mostly meson and the mixing parameters are small. Thus the  $f_0(1370)$  is not required from the point of view of mixing.

I show in Fig.2 (right) the probability distribution for the final mostly glueball state. It turns out that the experimental  $f_0(1710)$  comes out mostly glueball in agreement with the lattice calculation of mixing [2, 3] and other phenomenological analysis [9].

Let me now elaborate on the other scenario, namely assuming  $m_g > 1710$  MeV, i.e. I take 1750 MeV, close to the average value, as the unmixed glueball mass. As a result of Cardano's analysis, I know that the only way to get that mass down to the  $f_0(1710)$  is by incorporating a heavier meson. In particular, I show in Fig. 3 (left) a three state mixing where the  $f_0(1710)$  is mostly glueball but the  $f_0(2020)$ , a not yet confirmed particle, must exist to get the experimental values for the lower masses. The corresponding probability curves, Fig. 3 (right), show that the glueball component is at the level of 70%. Most of the other 30 % is carried by the  $f_0(2020)$ . Besides the existence of the  $f_0(2020)$ , another feature of this scheme is that the  $f_0(1370)$ , whatever it be, is not required, and thus the error bars are really very small and the fit quite restrictive. Another characteristic of this weak mixing scheme in the spread distribution of the unmixed spectrum: the particles appear with an energy step of 150 – 200 MeV. Note that in this case a two state mixing scenario could




Figure 4: Eigenvalues of the mixing matrix for the glueball and two meson states. The masses and their errors are from PDG [10].

also be trivially constructed, but here the  $f_0(2020)$  would be unavoidable and the  $f_0(1500)$  would decouple from the mixing scheme.

Within the latter scenario also a strong mixing realization arises. As shown in Fig.4, I am able to get a good fit to the data starting from almost degenerate unmixed states. In this case the unmixed values are for two mesons at 1710 MeV and 1760 MeV and a glueball at 1750 MeV. The mixing parameters between the almost degenerate light meson and glueball is large  $b = 210$  MeV, while the others are normal  $a = 100$  MeV,  $c = 70$  MeV. On the other hand this strong mixing affects the  $f_0(1710)$  strongly, which now contains very little glueball,  $\sim 17\%$ , while increasing dramatically the glueball content of the  $f_0(1500)$ ,  $\sim 41\%$ , and some that of the  $f_0(2020)$ ,  $\sim 42\%$ , see Fig 5. In this strong mixing scenario the glueball is a relic of the original glueball state and therefore it will be difficult to single out glueball properties. Again no need for the  $f_0(1370)$  in the fit.




Figure 5: The probability distributions as a function of  $b$  of the mixed states, for mixing parameters  $a = 100$  MeV,  $c = 70$  MeV. The notation follows the unmixed states: the full line represents the glueball (1750 MeV), the dashed line represents the meson (1710 MeV), the dotted line represents the meson (1760) MeV. The left figure represents the probability distribution of the  $f_0(2020)$ , the middle that of the  $f_0(1710)$ , and the right figure that of the  $f_0(1500)$ .

I might summarize at this stage my findings by stating that accepting a mass value for a glueball as obtained from lattice QCD for the unmixed glueball state and allowing for mixing under the strict scrutiny of the  $1/N_c$  expansion, two scenarios appear as dictated by the shape of the unmixed spectrum: i) a weak mixing scenario, whose consequence is that the  $f_0(1710)$  is an almost pure glueball state; ii) a strong mixing scenario in which the  $f_0(1710)$  has almost no glueball component, while the  $f_0(1500)$  and the  $f_0(2020)$  have about 40% glueball component. The role played by the scalar meson spectrum in the realization of mixing is fundamental. A detailed lattice QCD description of this spectrum would determine the mixing scenario. Within the weak mixing scenario it is very important the value of the mass of the unmixed glueball state. If it is below the  $f_0(1710)$  the mixing proceeds via the  $f_0(1500)$  and the  $f_0(1370)$  and it is not very constrained due to the large indeterminacy in the latter. If that mass is above 1710 MeV the  $f_0(2020)$  is required to bring the mass value down, while the  $f_0(1370)$  is unnecessary.

## 4 Conclusions

In this paper I take a very pragmatic point of view. I assume that the lowest found lattice QCD glueball state is associated with real physical states within the  $f_0$  spectrum and try to establish the connection by implementing phenomenologically mixing with the nearby scalar mesons. In order to estimate the mixings I use the splitting pattern of the  $f_0$  spectrum and  $1/N_c$  arguments.

I find two possible weak mixing scenarios which associate the lattice glueball mostly with the  $f_0(1710)$ . In this case the exact value of the unmixed glueball state is fundamental to determine the physical realization of mixing, being the  $f_0(1710)$  the dominant scale. If the unmixed glueball mass is below that value, the  $f_0(1370)$  should exist and may play a relevant role. If the mass is above, at least the  $f_0(2020)$  is required to enter the mixing to bring down the unmixed mass to the physical mass. The scalar spectrum in both cases is loose, with an energy step between 150 – 200 MeV.

I also find a strong mixing scenario which leads to the  $f_0(1500)$  and the  $f_0(2020)$  having a strong glueball component, while the  $f_0(1710)$  is mostly mesonic in character. In this case the unmixed scalar spectrum presents an almost degeneracy around the unmixed glueball mass.

The main conclusion of this calculation is that given the fact that mixings are difficult to calculate in lattice QCD, a good knowledge of the scalar meson spectrum together with phenomenology will clarify the glueball constituency of the physical  $f_0$ 's in full QCD. Once the scalar spectrum is known the  $f_0$  spectrum and the arguments about the mixing used in this communication will fix quite strongly the mixing parameters.

Giving the above analysis I find that the  $f_0(1710)$  being mostly glueball is the most natural scenario. If it is accompanied, besides the  $f_0(1500)$ , by the  $f_0(1370)$  or the  $f_0(2020)$  in the mixing scenario is a matter of experimental determination. It is clear that for this purpose not only masses, but also decays will be needed to determine the main properties of the glueball component. In here I have presented a guide of possible mixing schemes as characterized by the structure of the spectrum.

## Acknowledgments

Correspondence with Jaume Carbonell and Vincent Mathieu is gratefully acknowledged. This work was supported by the MINECO under contract FPA2013-47443-C2-1-P, by GVA-PROMETEOII/–2014/066, and by CPAN(CSD-00042).

## References

- [1] V. Mathieu, N. Kochelev and V. Vento, *Int. J. Mod. Phys. E* **18** (2009) 1 [arXiv:0810.4453 [hep-ph]].
- [2] W. J. Lee and D. Weingarten, *Phys. Rev. D* **61** (2000) 014015 [hep-lat/9910008].
- [3] A. Vaccarino and D. Weingarten, *Phys. Rev. D* **60** (1999) 114501 [hep-lat/9910007].
- [4] C. J. Morningstar and M. J. Peardon, *Phys. Rev. D* **60** (1999) 034509 [hep-lat/9901004].
- [5] Y. Chen, A. Alexandru, S. J. Dong, T. Draper, I. Horvath, F. X. Lee, K. F. Liu and N. Mathur *et al.*, *Phys. Rev. D* **73** (2006) 014516 [hep-lat/0510074].
- [6] E. Gregory, A. Irving, B. Lucini, C. McNeile, A. Rago, C. Richards and E. Rinaldi, *JHEP* **1210** (2012) 170 [arXiv:1208.1858 [hep-lat]].
- [7] C. Amsler and F. E. Close, *Phys. Rev. D* **53** (1996) 295 [hep-ph/9507326].
- [8] D. Barberis *et al.* [WA102 Collaboration], *Phys. Lett. B* **397** (1997) 339.
- [9] H. Y. Cheng, *Int. J. Mod. Phys. A* **24** (2009) 3392 [arXiv:0901.0741 [hep-ph]].
- [10] K. A. Olive *et al.* [Particle Data Group Collaboration], *Chin. Phys. C* **38** (2014) 090001.